

# Illustrating Clinic



Immediately after  
Surgery



5 days  
postoperative



14 days  
postoperative

# Illustrating Clinic



Patient Nguyen Thi L 40years old. Admission 15/12/2014

Patient with wound at finger 2,3

# Illustrating Clinic



Oblique skin flap of back finger 3,4 cover soft defect  
Skin thickness graft in donor site

# Illustrating Clinic



Patient Nguyen Van V 19 years old with amputated wound of fingertips 2,3,4 at second hour

# Illustrating Clinic



V-Y flap cover soft defect

# Illustrating Clinic



10 days postoperative

# Illustrating Clinic



Patient Nguyen Van V 32years old. Admission 06/2/2015

Wound with whole defect of palmar second phalanx of finger 1 at fourth

Using fasciocutaneous pedicle flap in back of second phalanx of finger 2 with size: 3x4 cm

# Illustrating Clinic



# Illustrating Clinic



7 days postoperative, entire survival of flap,  
location of skin graft was good

# Illustrating Clinic



20 days postoperative, skin color grafting is similar to the surrounding skin, good life of skin flap, normal movement of thumb

## 4. Conclusion

The results demonstrate once again that all skin at the wrist, hand and fingers is a reservoir for the flexible design of fasciocutaneous pedicle flap .

Emergency surgery not only covers tendon, bones exposed, limits infection, but also contributes to rapid wound cleaning, increase the feed source, promote the process of healing, especially facilitates the rehabilitation of the function of the wrist, hand and fingers and shortens the time, minimizing the cost of treatment.

**Thanks for your attention !**

